


AMENDMENTS TO THE CLAIMS

The following listing of claims replaces all prior versions and listings of claims for this application.

1. (original) A process for the oxidation of an unsaturated alcohol of formula




wherein:

R¹ represents a hydrogen atom, a C₁ to C₂₀ linear, branched or cyclic saturated or unsaturated hydrocarbon group, said hydrocarbon group optionally being substituted and also optionally including one or two oxygen or nitrogen atoms;

R² represents a C₂ to C₂₀ linear, branched or cyclic alkenyl, alkandienyl or alkantrienyl hydrocarbon group, said hydrocarbon group optionally being substituted and also optionally including one or two oxygen or nitrogen atoms; or said R¹ and R² optionally may be bonded together to form an unsaturated ring having 5 to 20 carbon atoms, said ring optionally being substituted and also optionally including one or two oxygen or nitrogen atoms; wherein the optional substituents of R¹, R² and of the ring which said R¹ and R² together may optionally form, are C₁ to C₁₅ linear, branched or cyclic alkyl, alkenyl or aromatic groups;

into a corresponding unsaturated aldehyde or ketone of formula




wherein R¹ and R² are as defined in formula (I);

wherein the oxidation is performed by a hypochlorite salt of formula



and in the presence of a catalytic amount of a N-(2,2,6,6-tetraalkyl-4-piperidiny-N-oxyl)-2-amino-1,3,5-triazine compound.

2. (original) The process of claim 1, wherein the unsaturated alcohol is a compound of formula (I),




wherein:

R¹ represents a hydrogen atom;

R² represents a C₅ to C₁₅ linear, branched or cyclic alkenyl or alkandienyl hydrocarbon group, that is optionally substituted; or R² represents a C₇ to C₁₅ linear, branched or cyclic alkantrienyl hydrocarbon group that is optionally substituted; and the optional substituents of R² are C₁ to C₈ linear, branched or cyclic alkyl, alkenyl or aromatic groups.

3. (original) The process of claim 1, characterized in that the N-(2,2,6,6-tetraalkyl-4-piperidiny-N-oxyl)-2-amino-1,3,5-triazine is of formula



wherein z represents an integer from 1 to 20;


R³ represents, simultaneously or independently, a hydrogen atom or an oxyl radical (O•), with the proviso that at least one R³ group is an oxyl radical;

X represents an oxygen atom or a $-\text{NR}^4-$ group;

R⁴ represents, simultaneously or independently, a hydrogen atom, a 2,2,6,6-tetramethyl-4-piperidinyl group, a 2,2,6,6-tetramethyl-4-piperidinyl-N-oxyl radical group or a C₁ to C₁₅ linear, branched or cyclic saturated or unsaturated hydrocarbon group, said hydrocarbon group optionally including one or two oxygen or nitrogen atoms; or two R⁴ groups, bonded to the same nitrogen atom, may be bonded together to form a heterocycle having 5 to 7 members and which may contain an oxygen atom;

R⁵ represents, simultaneously or independently, a hydrogen atom or a NR⁶₂ group;


R⁶ represents, simultaneously or independently, a hydrogen atom, a C₁ to C₂₀ linear, branched or cyclic saturated or unsaturated hydrocarbon group, a 2,2,6,6-tetramethyl-4-piperidinyl-N-oxyl radical group, a 2,2,6,6-tetramethyl-4-piperidinyl group or a group of formula



R⁷ representing, simultaneously or independently, a hydrogen atom, a C₁ to C₁₂ linear or branched alkyl group, a 2,2,6,6-tetramethyl-4-piperidinyl-N-oxyl radical group or a 2,2,6,6-tetramethyl-4-piperidinyl group; and

Y represents, simultaneously or independently, a C₂ to C₂₀ linear, branched or cyclic alkylene group optionally including one or two oxygen or nitrogen atoms.

4. (original) The process of claim 3, wherein the N-(2,2,6,6-tetraalkyl-4-piperidinyl-N-oxyl)-2-amino-1,3,5-triazine compound is a polymeric or oligomeric compound of formula



wherein z represents an integer from 2 to 10;

m represents an integer from 2 to 12;

R³ is as defined in claim 3;

R⁴ represents, simultaneously or independently, a hydrogen atom, a 2,2,6,6-tetramethyl-4-piperidiny-N-oxyl radical group, a 2,2,6,6-tetramethyl-4-piperidiny group or a C₁ to C₁₀ linear or branched alkyl or alkenyl group; or two R⁴ groups, bonded to the same nitrogen atom, may be bonded together to form a heterocycle having 6 members and which may contain an oxygen atom; and

R⁸ represents, simultaneously or independently, a hydrogen atom, a C₁ to C₁₀ linear or branched alkyl or alkenyl group, a 2,2,6,6-tetramethyl-4-piperidiny-N-oxyl radical group, a 2,2,6,6-tetramethyl-4-piperidiny group or a group of formula (V).

5. (original) The process of claim 4, wherein the N-(2,2,6,6-tetraalkyl-4-piperidiny-N-oxyl)-2-amino-1,3,5-triazine derivative is a N-oxyl derivative of the polymers having the CAS Registry Numbers 71878-19-8 or 192268-64-7.

6. (original) The process of claim 1, wherein the hypochlorite salt is selected from the group consisting of NaOCl, KOCl and Ca(OCl)₂.

7. (original) The process of claim 1, wherein one of a bromide salt of formula M'Br or a bicarbonate of formula M'HCO₃ is added to the process, wherein M' is an alkaline metal.

8. (original) The process of claim 1, wherein a bromide salt of KBr or NaBr is added to the process.

9. (original) The process of claim 1, wherein a bicarbonate of KHCO₃ or NaHCO₃ is added to the process.

10. (original) The process of claim 1, wherein the catalyst has a concentration ranging from 0.02 to 0.15 molar equivalents relative to the amount of alcohol.

11. (original) The process of claim 1, wherein the hypochlorite salt is added to the reaction mixture in an amount of between 0.9 and 2.5 molar equivalents relative to the amount of alcohol.

12. (currently amended by being renumbered) The process of claim 1, which is carried out at a temperature ranging between 0°C and 60°C

13. (currently amended by being renumbered) The process of claim 1, conducted in a solvent.

14. (currently amended by being renumbered) The process of claim 1, conducted in the absence of a solvent.